

Exam 4a

Chem 1121

Summer 2008

Name: _____

Take a deep breath, and relax! First, answer the questions you know how to do and then work on the more difficult problems. Don't forget to show all your work, so I can give you as much credit as possible.

Good Luck!

Andy

Q1. The reaction between stomach acid (HCl) and tums (CaCO₃) is given by the following **unbalanced** chemical equation:



a) [10 pts.] Balance the equation using the *lowest* set of whole number coefficients.

b) [7 pts.] Using the conversion factor method, calculate how many moles of CO₂ will be produced when 0.24 mol of HCl is neutralized.

c) [10 pts.] Using the conversion factor method, calculate the mass of CO₂ that will be produced from 0.450 g of CaCO₃

Q2. Convert the following masses to moles:
a) [5 pts.] 0.35 g of CH_2O , formaldehyde

b) [5 pts.] 12.0 g of $\text{C}_6\text{H}_{12}\text{O}_6$, glucose.

Q3. [5 pts.] What mass would 0.50 mol of water, H_2O , weigh?

Q4. *Use the conversion factor method for both of the problems.*
[5 pts.] How many moles of glucose are contained in 0.200 L of a solution that is 0.34 M?

[5 pts.] What volume of 0.34 M glucose solution contains 1.00 mol glucose?

Q5. [8 pts.] Sketch a diagram of a toricelli barometer, and explain how it can be used to measure atmospheric pressure.

Q6. [6 pts.] The total pressure of a mixture of three gases is 452 mmHg. If two of the three gases have partial pressures of 124 mmHg and 201 mmHg, then what must be the partial pressure of the third gas?

Q7. [10 pts.] Match the gas laws:

- | | |
|---------------------|--------------------|
| 1) Boyle's Law | a) $V \propto T$ |
| 2) Charles' Law | b) $pV = nRT$ |
| 3) Avogadro's Law | c) $p \propto T$ |
| 4) Gay Lussac's Law | d) $V \propto n$ |
| 5) Ideal Gas Law | e) $P \propto 1/V$ |

Q8. [7 pts.] A balloon of gas with a volume of 23.0 L at a pressure of 742 mmHg is squeezed so that its pressure becomes 921 mmHg. What will its new volume be?

Q9. [7 pts.] What volume will 0.15 mol of a gas occupy if its temperature is 23 °C, and its pressure is 0.45 atm?

Q10. [5 pts.] **Fill in the blanks.** The pressure of gas above a liquid is known as the _____ pressure. At the boiling point of a liquid, this pressure is equal to _____.

Q11. [5 pts.] Using the kinetic theory of gases (the idea that gases are composed of a large number of tiny particles in a state of constant chaotic motion) explain what is responsible for the **pressure** of a gas?

BONUS QUESTION:

What is the name given to a solution that has the *maximum* amount of solute dissolved in a given amount of solvent.

Periodic Table of the Elements

IA 1	IIA 2											IIIA 13	IVA 14	VA 15	VIA 16	VIIA 17	VIIIA 18																
1 H 1.01																		2 He 4.00															
3 Li 6.94	4 Be 9.01											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18																
11 Na 22.99	12 Mg 24.31	13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.07	17 Cl 35.45	18 Ar 39.95	19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.87	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.39	31 Ga 69.72	32 Ge 72.61	33 As 74.92160	34 Se 78.96	35 Br 79.90	36 Kr 83.80								
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc [98]	44 Ru 101.07	45 Rh 102.91	46 Pd 106.42	47 Ag 107.87	48 Cd 112.41	49 In 114.82	50 Sn 118.71	51 Sb 121.76	52 Te 127.60	53 I 126.90	54 Xe 131.29	55 Cs 132.91	56 Ba* 137.33	57 La 138.91	58 Ce 140.12	59 Pr 140.91	60 Nd 144.24	61 Pm [145]	62 Sm 150.36	63 Eu 151.96	64 Gd 157.25	65 Tb 158.93	66 Dy 162.50	67 Ho 164.93	68 Er 167.26	69 Tm 168.93	70 Yb 173.04
87 Fr [223]	88 Ra** [226]	103 Lr [262]	104 Rf [261]	105 Db [262]	106 Sg [266]	107 Bh [264]	108 Hs [265]	109 Mt [268]	110 [269]	111 [272]	112 [277]	113 [285]	114 [289]	115 [293]	116 [297]	117 [290]	118 [294]	89 Ac [227]	90 Th 232.04	91 Pa 231.04	92 U 238.03	93 Np [237]	94 Pu [244]	95 Am [243]	96 Cm [247]	97 Bk [247]	98 Cf [251]	99 Es [252]	100 Fm [257]	101 Md [258]	102 No [259]		

$$1 \text{ atm} = 760 \text{ mmHg} = 760 \text{ torr} = 101,325 \text{ Pa}$$

$$pV = nRT$$

$$P_1V_1 = P_2V_2$$

$$V_1/T_1 = V_2/T_2$$

$$P_1/T_1 = P_2/V_2$$

$$R = 0.08206 \text{ L} \cdot \text{atm}/\text{mol} \cdot \text{K}$$

$$T(\text{K}) = t(^{\circ}\text{C}) + 273$$